

least, our results could be positively fundamental to in-depth studies about officially using PHA as a method to treat infection in the future.

References

- Amini F., Semnani D., Karbasi S., Banitaba S.N., 2018, A novel bilayer drug-loaded wound dressing of PVDF and PHB/Chitosan nanofibers applicable for post-surgical ulcers. *International Journal of Polymeric Material and Polymeric Biomaterials*, 68, 772–777.
- Chen G.Q., Jiang X.R., 2018, Engineering microorganisms for improving polyhydroxyalkanoate biosynthesis, *Current Opinion in Biotechnology*, 53, 20–5.
- Chun-Ji Ng., Xin-Fang L., Norliana M., Siti K. A., Yusof K., 2014, Kamsiah J., Reprint of "heated vegetable oils and cardiovascular disease risk factors", *Vascular Pharmacology*, 62(1), 38-46.
- Favaro L., Basaglia M., Casella S., 2019, Improving polyhydroxyalkanoate production from inexpensive carbon sources by genetic approaches: a review, *Biofuels, Bioproducts and Biorefining homepage*, 13(1), 208–27.
- Griffin G., 1993, *Chemistry and Technology of Biodegradable Polymers*, Chapman and Hall, London, UK, 1994th edition.
- Hageman G., Kikken R., Hoor F. T., Kleinjans J., 1988, Assessment of mutagenic activity of repeatedly used deep-frying fats, *Mutation Research*, 204(4): 593-604.
- Jiang X.R., Wang H., Shen R., Chen G.Q., 2015, Engineering the bacterial shapes for enhanced inclusion bodies accumulation, *Metabolic Engineering*, 29, 227–37.
- Kheang L. S., May. C. Y., Foon C. S., Ngan M. A., 2006, Recovery and Conversion of Palm Olein-Derived Used Frying Oil to Methyl Esters for Biodiesels, *Journal of Oil Palm Research*, 18, 247-252.
-
- Kundrat V., Cernekova N., Kovalcik A., Enev V., Marova I., 2019, Drug Release Kinetics of Electrospun PHB Meshes, *Materials (Basel)*, 12,1924.
- Li T., Ye J., Shen R., Zong Y., Zhao X., Lou C., Chen G.Q., 2016, Semirational approach for ultrahigh poly(3-hydroxybutyrate) accumulation in *Escherichia coli* by combining one-step library construction and high-throughput screening, *ACS Synthetic Biology*, 5(11), 1308–17.
- Ling C., Qiao G.Q., Shuai B.W., Olavarria K., Yin J., Xiang R.J., Song K.N., Shen Y.H., Guo Y., Chen G.Q., 2018, Engineering NADH/NAD⁺ ratio in *Halomonas bluephagenesis* for enhanced production of polyhydroxyalkanoates (PHA), *Metabolic Engineering*, 49,275–86.
- Mannu A., Garroni S., Porrás J. I., Mele A., 2020, Available Technologies and Materials for Waste Cooking Oil Recycling, *Processes*, 8(3), 366.
- Naveen N., Kumar R., Balaji S., Uma T., Natrajan T., Sehgal P., 2010, Synthesis of Nonwoven Nanofibers by Electrospinning—A Promising Biomaterial for Tissue Engineering and Drug Delivery, *Advanced Engineering Materials*, 12, B380–B387.
- Nikel P.I., Chavarría M., Danchin A., de Lorenzo V., 2016, From dirt to industrial applications: *Pseudomonas putida* as a synthetic biology chassis for hosting harsh biochemical reactions, *Current Opinion in Chemical Biology*, 34, 20–9.
- Orjuela A., Clark J., 2020, Green chemicals from used cooking oils: Trends, challenges, and opportunities, *Current Opinion in Green and Sustainable Chemistry*, 26: 100369.
- Poblete-Castro I., Binger D., Oehlert R., Rohde M., 2014, Comparison of mcl-Poly(3-hydroxyalkanoates) synthesis by different *Pseudomonas putida* strains from crude glycerol: citrate accumulates at high titer under PHA-producing conditions, *BMC Biotechnology*, 14, 962.
- Spaans S., Weusthuis R., Van Der Oost J., Kengen S., 2015, NADPH-generating systems in bacteria and archaea, *Frontiers in Microbiology*, 6, 742.
- Sudarsan S., Dethlefsen S., Blank L.M., Siemann-Herzberg M., Schmid A., 2014, The functional structure of central carbon metabolism in *Pseudomonas putida* KT2440, *Applied and Environmental Microbiology*, 80(17), 5292–303.
- Tan D., Wu Q., Chen J.C., Chen G.Q., 2014, Engineering *Halomonas* TD01 for the low-cost production of polyhydroxyalkanoates, *Metabolic Engineering*, 26, 34–47.
- Zhuang Q., Wang Q., Liang Q., Qi Q., 2014, Synthesis of polyhydroxyalkanoates from glucose that contain medium-chain-length monomers via the reversed fatty acid beta-oxidation cycle in *Escherichia coli*, *Metabolic Engineering*, 24, 78–86.